

The Relationship between Physical Activity and Body Mass Index: Issues in Model Specification

Gizachew Tiruneh, Ph. D., Department of Political Science, University of Central Arkansas

[A Revised version of this paper is published by the *International Journal on Disability and Human Development*, 2009, Volume 8, Number 3, pp. 267-275]

Abstract

The inverse relationship between physical activity and body fat or weight is well-known. What are not clearly understood, however, are the statistical models that best describe the relationship between the two variables. While many researchers employ the inverse linear estimator to explain the effect of physical activity on body fat or weight, others utilize the inverse logarithmic model. This study finds that the inverse logarithmic model is more accurate in describing the relationship between the two variables than is the inverse linear model. In addition, by specifying physical activity as a continuum of inactivity and activity, this study finds that although the inverse logarithmic method can be used in some cases, the inverse logistic model is best suited in describing the relationship between physical activity and body weight.


Introduction

Researchers, health specialists, and public officials have long known that an increased level of exercise is negatively related with body fat or weight. Numerous empirical studies have confirmed such a relationship (Johnson, Burke, & Mayer, 1956; Bullen, Reed, & Mayer, 1964; Tremblay, Despres, Leblanc, Lynn, Ferris, & Stephens et al., 1990; Tudor-Locke, Ainsworth, Thompson, Addy, & Jones, 2001; Thompson, Rakow, & Perdue, 2004; Krum, Dessieux, Andrews, & Thompson, 2006). The relationship between physical activity and body fat or weight is derived from the assumption that a normal-weight person's energy intake is equal or nearly equal to his/her energy expenditure (Pennington, 1953; Johnson et al., 1956; Bunyard, Kartzel, Busby-Whitehead, Wu, & Goldberg, 1998). That is, a person becomes overweight or obese if his/her energy intake is greater than his/her energy expenditure, and one way of maintaining the energy balance is by getting rid of the extra calories by performing physical activity (Esparza, Fox, Harper, Bennett, Schultz, & Valencia et al., 2000). What are not clearly understood, however, are the statistical models that best describe the connection between physical activity and body fat or weight (Dwyer, Hosmer, Hosmer, Venn, Blizzard, & Granger et al., 2007). For instance, Hemmingsson & Ekelund (2007) find that the association between physical activity and body mass index (BMI) is stronger in obese individuals than in non-obese persons. Their findings led them to question the conventional inverse-linear effect of physical activity on BMI. Not inconsistent with Hemmingsson & Ekelund's (2007) finding, Dwyer et al. (2007) show that the shape of the relationship between physical activity and BMI is inverse logarithmic. The insight that an inverse logarithmic model describes the physical activity and body fat or weight linkage,

however, is not new. Indeed, several decades ago, Mayer, Roy, & Mitra (1956; see also Bender & Bender, 1976), for instance, demonstrated an empirical pattern of a decreasing effect of physical activity on body weight. Subsequently, several other researchers have obtained similar results (Klem, Wing, McGuire, Seagle, & Hill, 1997; Haapanen, Miilunpalo, Pasanen, Oja, & Vuori, 1997; Tate, Jeffery, Sherwood, & Wing, 2007). Such findings suggest the presence of a threshold of physical activity, which occurs when a person moves from an overweight to a normal-weight status.

Given the usage of both the inverse linear and logarithmic estimators in previous research, the question becomes, which of the two models is more accurate in describing the relationship between physical activity and body fat or weight? In this study, using a cross-sectional study and a sample including obese, overweight, and normal-weight individuals, we find that although the inverse linear model provides significant results between the two variables, more variance in body weight is explained by employing the inverse logarithmic model. Hypothetical inverse linear and logarithmic curves are shown in Fig. 1.

Fig. 1: Hypothetical Inverse Linear and Logarithmic Curves



As will be made clear below, the inverse logarithmic model is very useful and can be employed in describing some aspects of the relationship between physical activity and body weight. However, previous research findings on the relationship between physical inactivity, as opposed to physical activity, and body fat or weight (Mayor et al., 1956; Johnson et al., 1956; Bullen et al., 1964) lead us to conceive that the best model for testing the relationship between physical inactivity/activity and body fat or weight among obese, overweight, and normal-weight persons may not be inverse logarithmic but logistic or probit. A key point for our thesis in the inactivity-related studies is the argument that a threshold of physical activity exists between sedentary and non-sedentary life styles. This threshold is distinct from the threshold that is believed to exist after an overweight individual or group achieves a normal-weight status. Mayor et al. (1956), for

instance, have empirically found that a “sedentary” range or threshold of physical activity is associated with obesity. Mayor et al.’s (1956) threshold argument has been supported by Schoeller’s (1998) empirical findings. Consistent with Mayer et al.’s (1956) and Schoeller’s (1998) threshold arguments, other researchers have provided various minimum thresholds of physical activity necessary to prevent a sedentary life style, including energy spent on steps per day (Hatano, 1993; Tudor-Locke et al., 2001; Thompson et al., 2004), time spent on moderate or rigorous exercise (Schoeller, Shay, & Kushner, 1997), and weekly caloric-expenditure (Jeffery, Wing, Sherwood, & Tate, 2003).

Given the foregoing, we argue that the pattern of the relationship between physical inactivity/activity and body fat or weight, with its upper and lower thresholds, would be best described by the logistic or probit model. A hypothetical inverse logistic curve is shown in Fig. 2. The description of physical activity (or lack there of) and body fat or weight by the logistic or probit model, however, requires a modification in the specification of the two variables. For instance, we could specify physical inactivity and activity as one variable varying in a continuum. More often than that, researchers treat physical inactivity distinct from or as an aspect of physical activity. Physical inactivity is specified as a decreasing level of activity (Mayer et al., 1956; Schoeller, 1998; Esparza et al., 2000) or a variable that varies from little to high level of inactivity (Treath, Hou, Young, & Maynard, 2005; Salmon, Bauman, Crawford, Timperio, & Own, 2000; Hemmingsson & Ekelund, 2007; Metallinos-Katsaras, Freedson, Fulton, & Sherry, 2007). It is also sometimes assumed to take a value of zero (or considered as a non-varying variable). We believe that the description of physical inactivity as a decreasing

level of activity is compatible with our specification of the inactivity-activity continuum. In other words, we prefer the description of inactivity as time lost in not exercising. Thus, time lost in things like television viewing and video-game playing should be considered only as aspects of inactivity. Similarly, the use of the logistic or probit model suggests the specification of body fat or weight as a variable that is continuous within and across the obese-overweight-normal weight categories. Thus, whereas physical inactivity would correspond to the obese, lesser and greater activities would fit the overweight and the normal-weight groups, respectively. And whereas the upper-flat part of the logistic or probit curve would describe obesity, the middle and the lower-flat portions would, respectively, represent an on-going weight loss and the maintenance of a healthy weight. Indeed, the empirical evidence in this study seems to provide support for our thesis.

In summary, this study will show that while the inverse logarithmic model is more accurate in testing the relationship between physical activity and body weight than is the inverse linear estimator, the logistic model is best suited in describing the linkage between the two variables.


Method

Research Design and Specification of Variables

This study employs a cross-sectional research design and ordinary least square (OLS) and Logistic regressions to test the impact of physical activity on BMI. BMI is considered to gauge 'bodyweight' as opposed to 'body fat', but it is argued that there is a correlation between the two variables (Ruderman, Schneider, & Berchtold, 1981; Cameroon & Getz, 1997). We calculated the BMI of the participants (the dependent

variable) using the standard formula, $\text{weight} / \text{height}^2$. For the logistic models, the BMI

Fig. 2: Hypothetical Inverse Logistic or Probit Curve



data are specified as follows: we assign 1 to individuals who are obese or overweight and a 0 to those who are neither obese nor overweight. Physical activity is measured by two variables: the frequency of exercise (the number of hours that a person works out in a week) and the duration of exercise (how long a person has been doing his/her weekly workout). Yet, physical activity is not the only factor that affects BMI. A host of factors including occupational activity (King, Fitzhugh, Bassett, McLaughlin, Strath, Swartz, et al., 2001), stress (Roberts, Troop, Connan, Treasure, & Campbell, 2007), smoking and socioeconomic status (Haapanen et al., 1997), drinking (Lahti-Koski, Pietinen,

Heliövaara, & Vartiainen, 2002), diet (Mitchell, 1952; Warnold, Carlgren, & Krotkiewski, 1978; Klem et al., 1997; Lahti-Koski et al., 2002; Metallinos-Katsaras et al., 2007), being in school or not and seasonal variations (Rowlands & Hughes, 2006), physical fitness (Bovett, Auguste, & Burdette, 2007), personal health (Payn, Pfeiffer, Hutto, Vena, LaMonte, & Blair et al., 2008), and genetics (Pennington, 1953; Lahti-Koski et al., 2002) seem to impact people's weight. The purpose of this study is not, however, the full specification of the variables that explain BMI. It is rather the specification of the models that describe BMI. Nevertheless, this study will control for the intensity of physical activity, age, and gender of the participants.

While some find that vigorous exercise is key to lower body fat or weight (Tremblay et al., 1990; Schoeller et al., 1997; Haapanen et al., 1997; Jeffery et al., 2003; Ruiz, Rizzo, Hurtig-Wennlof, Ortega, Warnberg, & Sjostrom, 2006; Tate et al., 2007; Metallinos-Katsaras et al., 2007), others show that moderate or light intensity is more important in achieving or maintaining a healthy body (Schoeller et al., 1997; Hills & Byrne, 1998; Treath et al., 2005). We classify the intensity of physical activity into low, moderate, and high categories. The low-intensity variable, which we expect to have the least benefits in affecting BMI, will be used as a baseline. Moreover, it has been argued that individuals' age and gender affect BMI. As people age, their weights tend to increase (Haapanen et al., 1997; Yanai, Kon, Kumasaka, & Kawano, 1997). Inactivity is also found to be positively related with body fat among boys but not among girls (Treath et al., 2005). Moreover, Cameron & Getz (1997) have found that adolescent females are more prone to obesity than males. Lastly, Weir, Ayers, Jackson, Rossum, Poston, & Foreyt (2001) have shown that gender is a predictor of long-term weight control.

Procedure and Data Collection

The participants in this study are undergraduate students at several U.S. universities. We collected two sets of data. For the first set (N=57), we administered a questionnaire to students at the University of Z. To make the data as nationally representative as possible, we collected a second set of data by administering an email-based questionnaire at 15 randomly selected universities by accessing the names of students through university directories. These universities are Rutgers, North Carolina, Florida, Auburn, Oklahoma, Texas, Oregon, Washington, Idaho, Utah, Colorado, South Dakota, Iowa, Ohio, and Memphis. The questionnaire was sent to about 25 students at each university. We received a total of 52 responses, a response rate of about five percent. The same questionnaire was used in both sets of data. The questionnaire asks students for their weight, height, sex, age, frequency, intensity, and duration of physical activity. The mean age of the combined data (N=107), which is skewed towards older continuing-education students, is 22.6. There are 53 male and 54 female students in the combined sample data. Of these students, 17 are obese, 15 are overweight, and 75 have weights in the normal range. The means for the duration of exercise for the obese, overweight, and normal-weight groups are about 24, 59, and 89 months, respectively. And the mean weights for the obese, overweight, and normal-weight groups are 35.54, 27.07, and 21.61 lbs, respectively. We administered the questionnaire at University of Z in the Spring of 2007, and the email-based-questionnaire was conducted in the Summer and Fall of 2007.

Results

We begin, in Table 1, by analyzing the impact of physical activity as well as the control variables on BMI. Model 1 and Model 2 estimate the data for students at the University of Z. We log the physical activity variables, Duration and Frequency; these variables will be called Logged-Duration and Logged-Frequency hereafter. Model 1 and 2 show the impact of the un-logged and logged physical activity variables on BMI, respectively. The only variables that are statistically significant at the 0.05 level in both models are Duration and Logged-Duration. In Model 3 and 4, we analyze the data for the other 15 universities. Model 3 and 4 show the impact of the un-logged and logged physical activity variables on BMI, respectively. In Model 3, the Duration and Age variables are significant at the 0.05 levels. The rest of the variables are insignificant. Model 4 shows that both the Logged-Duration and the Age variables are significant at the 0.05 and 0.10 levels, respectively. The rest of the variables are insignificant. We combine the data for the University of Z and the other 15 universities in Models 5 and 6. The results are very similar to those obtained in Model 3 and 4: the Logged-Duration and Age variables are significant in both models; the Age variable is, however, significant at the 0.10 level in Model 6. The rest of the variables are insignificant in both models. In analysis not shown here, we interacted the Frequency and Duration variables, but we found that the impact of the multiplicative effect on BMI is insignificant.

In Table 2, we compare the merits of the inverse linear and logarithmic models in describing the relationship between physical activity and BMI. The Duration and Logged-Duration variables represent the inverse linear and logarithmic models,

Table 1: OLS Regression Estimates; Dependent Variable: BMI						
	Model 1	Model 2	Model 3	Model 4	Model 5	Model 6
	B	B	B	B	B	B
Intercept	24.11** (4.10)	27.13** (4.40)	13.59** (4.38)	21.11** (4.23)	19.85** (2.51)	23.97** (2.74)
Duration	-0.03** (0.01)		-0.05** (0.01)		-0.03** (0.01)	
Logged-Duration		-1.32** (0.46)		-1.69** (0.60)		-1.39** (0.35)
Frequency	-0.06 (0.21)		0.36 (0.34)		0.01 (0.17)	
Logged-Frequency		-0.42 (1.35)		2.62 (2.08)		0.96 (1.10)
Gender	0.38 (1.60)	0.57 (1.56)	1.28 (1.67)	1.40 (1.69)	0.89 (1.10)	1.00 (1.10)
Age	0.14 (0.19)	0.60 (0.18)	0.55** (0.17)	0.27** (0.12)	0.32** (0.11)	0.19* (0.10)
High Intensity	-0.54 (2.73)	-0.14 (2.79)	-0.63 (4.02)	0.03 (4.02)	-0.12 (2.19)	0.03 (2.23)
Moderate Intensity	-0.25 (1.80)	-0.29 (1.81)	-2.47 (1.84)	-3.20 (1.95)	-1.44 (1.23)	-1.62 (1.27)
N:	55	55	52	52	107	107
R ²	0.14	0.16	0.27	0.25	0.16	0.18
Note: **: p < 0.05; *: p < 0.10; Bs are unstandardized betas; standard errors in parentheses						

respectively. We do not use the Frequency variable here since, as shown in Table 1, it shows little or no impact on BMI. Model 1 and 2 show the impact of physical activity on BMI for the data we obtained from the University of Z. Both Duration and Logged-

Duration are significant and have negative signs, but the r-squared value of the latter (0.16) is higher than the former (0.12). In Models 3 and 4, we show the impact of Duration and Logged-Duration, respectively, on the BMI data of the college students at

	Model 1	Model 2	Model 3	Model 4	Model 5	Model 6
	B	B	B	B	B	B
Intercept	26.78** (1.00)	29.24** (1.56)	25.29** (1.14)	28.87** (2.17)	26.05** (0.75)	29.10** (1.27)
Duration	-0.03** (0.01)		-0.01 (0.01)		-0.02** (0.01)	
Logged-Duration		-1.30** (0.42)		-1.25** (0.55)		-1.29** (0.33)
N:	55	55	52	52	107	107
R	-0.35**	-0.39**	-0.18	-0.31*	-0.27**	-0.36**
R ²	0.12	0.16	0.03	0.09	0.07	0.13
Note: **: p < 0.05; *: p < 0.10; Bs are unstandardized betas; standard errors in parentheses						

the other 15 universities. Whereas Duration fails to have any effect on BMI in Model 3, Logged-Duration is statistically significant in Model 4. In addition, the r-squared value is higher for the Logged-Duration variable (0.09) than is for Duration (0.03). Lastly, in Model 5 and 6, we combine the data for University of Z and the other 15 universities to compare the impact of Duration and Logged-Duration on BMI, respectively. Both the

Duration and Logged-Duration variables are statistically significant and have negative signs, but the r-squared value is higher for the latter variable (0.13) than is for the former (0.07). In sum, the analyses in Table 2 consistently show that the inverse logarithmic model is more accurate, and thus appropriate, in describing the relationship between physical activity and BMI than is the inverse linear model.

We noted earlier, however, that a theoretically sound description of the relationship between physical inactivity-activity and BMI when using samples that include obese, overweight, and normal weight individuals is the logistic or probit model. We employ both OLS and logistic models in Table 3 to verify the foregoing assumption. First, we use the Duration variable to explain BMI and run OLS regressions by parsing the BMI data into obese, overweight, and normal groups. We do this because if we are to use the inverse logistic model, we will need to observe that the correlation between physical activity and the overweight group is higher than it is on each of the other two groups. In other words, we are specifically interested in seeing the strength and sign of the correlation between physical activity and the BMI of each group. Model 1 shows that the correlation between Duration and obese individuals' BMI is only 0.08 and the sign of the association is positive. In Model 2, we regress BMI on the Duration variable of the overweight group; the correlation is higher, 0.33, compared to the obese group, and the sign is negative. The Duration variable is not significant, however, and this may be partly a consequence of the overweight group's small sample size (N=15). In Model 3, we regress BMI on the Duration variable of the normal-weight group. Similar to the obese, the correlation is only 0.06 and the sign is positive. In other words, Models 1 through 3 suggest that the negative correlation between physical activity and BMI is due

mainly to the overweight group. For comparison, we show the results for the combined data (N=107) in Model 4.

	Model 1	Model 2	Model 3	Model 4	Model 5	Model 6
	B	B	B	B	B	B
Intercept	35.24** (1.64)	27.51** (0.52)	21.49** (0.33)	26.05** (0.75)	0.59 (0.49)	-1.01 (1.26)
Duration	0.01 (0.04)	-0.01 (0.01)	0.001 (0.003)	-0.02** (0.01)		
Logged-Duration					-0.43** (0.14)	-0.51** (0.16)
Logged-Frequency						0.50 (0.47)
Gender						1.05** (0.49)
Age						0.03 (0.05)
High Intensity						0.84 (0.89)
Moderate Intensity						-0.57 (0.57)
N:	17	15	75	107	107	107
R	0.08	-0.33	0.06	-0.27**		
R ²					0.08	0.16
Note: **: p < 0.05; *: p < 0.10; Bs are unstandardized betas; standard errors in parentheses						

Although the correlation value (-0.27) is statistically significant ($p < 0.05$) in the combined data, it is lower than the correlation value of the overweight group (-0.33) shown in Model 2. In other words, the rate of weight reduction for the overweight seems to be higher than the rates for the obese or the normal-weight groups. The implication of these results is that the best model for describing the obese-overweight-normal weight continuum is logistic or probit.

Given the foregoing, in Models 5 and 6, we use the logistic model to estimate the relationship between physical activity and BMI for the combined data ($N=107$). We first show the sole impact of the Logged-Duration variable on BMI in Model 5. The Logged-Duration variable is significant at the 0.05 level. More specifically, the model [$L = 0.59 - 0.43$ (Logged-Duration)] suggests that for one unit of increase in the Logged-Duration variable, the log of odds of being obese or overweight decreases by 0.43 units. This also means that for one unit of increase in the Logged-Duration variable, the odds ratio of an individual being obese or overweight decreases by 35 percent.


In addition, using the predicted model of the logistic regression, in Fig. 3 (in inverted S-shaped curve) we show the overall probability pattern of being obese or overweight for given Logged-Duration values. For instance, the probability of an undergraduate student being obese or weight when exercising for about 20 months (Logged-Duration = +3) is 0.33 or 33 percent. In contrast, if a student has not exercised (or has been inactive) for about 20 months (Logged-Duration = -3), the probability of such an individual being obese or overweight is 0.87 or 87 percent. As shown in Fig. 3, a whole range of values for the inverse logistic curve are fitted across the negative and positive X-axis, because we are able to estimate or predict the probabilities for negative Logged-Duration values.

In other words, the negative Logged-Duration values are derived from the model and only represent the opportunity cost of not exercising. We cannot, however, compare the merits of the logistic and logarithmic models merely by the variance of BMI explained by the physical activity variable; this is because the two models rely on dichotomous and continuous dependent variables, respectively. Thus, the choice of the logistic model over logarithmic should be based on theoretical considerations.

Moreover, we show the rates of change in the probability of being obese or overweight for given Logged-Duration values in Fig. 3 (the bottom curve). The rates of change in probability with respect to X or Logged-Duration values form an inverted bell-shaped like curve. This curve is actually the inverse or derivative of the inverse logistic curve. The inverted curve shows that the highest probability of decrease in weight in this particular set of sample (10.8 percent) occurs when the Logged-Duration value is 1.37 or when a student exercises for about 4 months. This figure also corresponds to a 0.50 or 50 percent probability of being obese or overweight, which happens to occur exactly at the middle of the inverse logistic curve.

In Model 6, we control for the Intensity, Gender, and Age variables. Because the logged-variables have shown higher variances in explaining BMI than have their unlogged counterparts, we use Logged-Duration and Logged-Frequency as measures of physical inactivity-activity in this model. The only variables that are statistically significant at the 0.05 level are Logged-Duration and Gender.

Fig. 3: Physical Activity and BMI



Discussion

One interesting finding of this study is that the duration of exercise is that matters in lowering or maintaining the BMI of undergraduate-college students. Such a finding is, however, not surprising since a student who has just started exercising and another who has been doing so for a longer period of time should not be expected to gain the same benefits in weight reduction or maintenance.

Far more surprising, however, is our finding that the intensity of physical activity does not matter in lowering or maintaining BMI. One has to be careful, however, when interpreting such a result. For instance, respondents in this study might not have

accurately described the intensity of their physical activities. Experimental studies should verify this finding.

This study also finds that the inverse logarithmic model is more accurate in describing the effect of physical activity on BMI than is the inverse linear estimator. Perhaps the most interesting observation and implication of this study is that the relationship between physical inactivity-activity and BMI may be best captured best by the logistic model. The mathematical implication of the logistic curve (as it would be the case with the logarithmic model) is that the level of the BMI of an individual could decrease, albeit infinitesimally, indefinitely. In practice, the leveling off of BMI at the lower end of a curve is interpreted to mean just the stabilization and maintenance of a healthy weight by an individual. While such an interpretation is close to the truth, the mathematical insight also seems to be valid: even an individual with a healthy BMI needs to maintain or slightly lower his/her level of weight (preferably by continuously, albeit slightly, increasing his/her level of physical activity) until he/she is too old or sick to exercise. It is known that even a person with a normal weight could vary his/her weight to his/her lower and upper limits of acceptable weight by about 40 lbs. If physical activity is not sustained, even an individual with a healthy weight could easily regain some or all of it (see also Schoeller et al., 1997; Tate et al., 2007).

Gender variation does not seem to have a major effect on BMI. This variable is statistically significant only in one of the logistic models. Indeed, there is no good theoretical reason to assume that males or females have inherent tendencies to become obese or overweight, and our result is also consistent with several other inconclusive findings. Finally, the Age variable has shown significance in some of the models. This

is not quite surprising since it is well-known that as people age, their level of physical activity decreases.

Conclusions

Ultimately, the use of statistical models should be guided by theoretical considerations and by the research questions raised. Theories of the link between physical activity and body fat or weight have long suggested the presence of an inverse relationship between the two variables. What has not been well clarified, however, is how the inverse linear, logarithmic, or logistic models describe the relationship between physical activity and body fat or weight. We found that the inverse logarithmic model is more accurate in describing the effect of physical activity on the BMI of obese, overweight, and normal-weight groups than is the inverse linear estimator. We also found some evidence that when a cross-sectional study includes obese, overweight, and normal weight individuals, the relationship between physical inactivity-activity and BMI could be best explained by employing the logistic model.

Our findings also have several implications. For instance, experimental studies could be conducted using the logistic or probit model to measure the impact of physical activity on a group of obese individuals over time. Such longitudinal studies would show that the pattern of weight reduction of the obese group would initially be very slow, followed by a faster speed, and ending with a very slow (stabilization) period. If a longitudinal study is, however, interested in the weight reduction of an overweight group (but not the obese), a better estimator will be the inverse logarithmic model. This model would show that faster weight reduction of the overweight group would be followed by the group's slower

weight-stabilization period. The inverse logarithmic model could also be used both in cross-sectional and longitudinal studies when a study consists of both overweight and normal-weight individuals. In such scenarios, the weight reductions of the latter group would be much slower than the former.

Clearly, future research should replicate the findings and implications of this study. In addition, the impact of intensity of physical activity should be verified by using more direct and objective measures of BMI such as accelerometers and pedometers. Future studies should also be done among all and different age groups, not just on college-age individuals.

References

- Bender, E. A. & Bender, A. D. (1976). Maintenance of weight loss in obese subjects. *British Journal of Preventive & Social Medicine*, 30, 50-65.
- Bovett, P., Auguste, R., & Burdette, H. (2007). Strong inverse association between physical fitness and overweight in adolescents: a large school-based survey. *International Journal of Behavioral Nutrition and Physical Activity*, 4, 1-8.
- Bullen, A. B., Reed, B. R., & Mayer, J. (1964). Physical activity of obese and nonobese adolescent girls appraised by motion picture sampling. *American Journal of Clinical Nutrition*, 14, 211-223.
- Bunyard, B. L., Kartzel, I. L., Busby-Whitehead, M. J., Wu, Z., & Goldberg, P. A. (1998). Energy requirements of middle-aged men are modifiable by physical activity. *American Journal of Clinical Nutrition*, 68, 1136-1142.
- Cameroon, N. & Getz, B. (1997). Sex differences in the prevalence of obesity in rural African adolescents. *International Journal of Obesity*, 21, 775-782.
- Dwyer, T., Hosmer, D., Hosmer, T., Venn, J. A., Blizzard, I. C., Granger, H. R. et al. (2007). The inverse relationship between number of steps per day and obesity in a population-based sample – the AusDiab Study. *International Journal of Obesity*, 31, 797-804.
- Esparza, J., Fox, C. Harper T.I., Bennett, H. P., Schultz, O.L., Valencia, E. M., & Ravussin, E. (2000). Daily energy expenditure in Mexican and USA Pima Indians: low physical activity as a possible cause of obesity. *International Journal of Obesity*, 24, 55-59.
- Gujarati, D. (1995). *Basic Econometrics*. New York: McCraw-Hill.
- Haapanen, N., Miilunpalo, S., Pasanen, M., Oja, P., & Vuori, I. (1997). Association between leisure time physical activity and 10-year body mass change among working-aged men and women. *International Journal of Obesity*, 21, 288-296.
- Hatano, Y. (1993). Use of the pedometer for promoting daily walking exercise. *International Council for Health, Physical Education and Recreational Journal*, 29, 4-8.
- Henningsson, E. & Ekelund, U. (2007). Is the association between physical activity and body mass index obesity dependent? *International Journal of Obesity*, 31, 663-668.
- Hills, P. A., & Byrne, M. N. (1998). Exercise prescription for weight management. *Proc. Nutr. Soc*, 57, 93-103.
- Jeffery, W. R., Wing, R. R., Sherwood, E. N., & Tate, F. D. (2003). Physical activity and weight loss: does prescribing higher physical activity goals improve outcome? *American Journal of Clinical Nutrition*, 78, 684-689.
- Johnson, L. M., Burke, S. B., & Mayer, J. (1956). Relative importance of inactivity and overeating in the energy balance of obese high school girls. *American Journal of Clinical Nutrition*, 4, 37-44.
- King, A. G., Fitzhugh, C. E., Bassett, R. D., McLaughlin, E. J. Strath, J. S., Swartz, M. A. et al. (2001). Relationship of leisure-time physical activity and occupational activity to the prevalence of obesity. *International Journal of Obesity*, 25, 606-612.
- Klem, L. M., Wing, R.R., McGuire, T. M., Seagle, M. H., & Hill, O. J. (1997). A descriptive study of individuals successful at long-term maintenance of substantial weight loss. *American Journal of Clinical Nutrition*, 66, 239-246.

- Krum, M. E., Dessieux, L. O., & Thompson, L. D. (2006). The relationship between daily steps and body compositions in postmenopausal women. *Journal of Women's Health, 15*, 202-210.
- Lahti-Koski, M., Pietinen, P., Heliövaara, M., & Vartiainen, E. (2002). Associations of body mass index and obesity with physical activity, food choices, alcohol intake, and smoking in the 1982-1997 FINRISK studies. *American Journal of Clinical Nutrition, 75*, 809-817.
- Mayer, J., Roy, P., & Mitra, P. K. (1956). Relation between caloric intake, body weight, and physical work. *American Journal of Clinical Nutrition, 4*, 169-175.
- Metallinos-Katsaras, S. E., Freedson, S. P., Fulton, E. J., & Sherry, B. (2007). The association between an objective measure of physical activity and weight status in preschoolers. *Obesity, 15*, 686-694.
- Mitchell, H. H. (1952). Overnutrition and obesity. *American Journal of Clinical Nutrition, 1*, 66-76.
- Payn, T., Pfeiffer, A. K., Hutto, B., Vena, E. J., LaMonte, J., Blair, N. S., & Hooker, P. S. (2008). Daily steps in midlife and older adults: relationship with demographic, self-rated health, and self-reported physical activity. *Research Quarterly for Exercise and Sport, 79*, 128-132.
- Pennington, W. A. (1953). An alternate approach to the problem of obesity. *American Journal of Clinical Research, 1*, 100-106.
- Roberts, C., Troop, N., Connan, F., Treasure, J., & Campbell, C. I. (2007). The effects of stress on body weight: biological and psychological predictors of change in BMI. *Obesity, 15*, 3045-3055.
- Rowlands, V. A. & Hughes, R. D. (2006). Variability of physical activity patterns by type of day and season in 8-10-year-old boys. *Research Quarterly for Exercise and Sport, 77*, 391-395.
- Ruderman, B. N., Schneider, H. S., & Berchtold, P. (1981). The "Metabolically-obese," normal-weight individual. *American Journal of Clinical Nutrition, 34*, 1617-1621.
- Ruiz, R. J., Rizzo, S. N., Hurtig-Wennlof, A., Ortega, B. F., Warnberg, J., & Sjostrom, M. (2006). Relations of total physical activity and intensity to fitness and fatness in children: the European youth heart study. *American Journal of Clinical Nutrition, 84*, 299-303.
- Salmon, J., Bauman, A., Crawford, D., Timperio, A., & Owen, N. (2000). The association between television viewing and overweight among Australian adults participating in varying levels of leisure-time physical activity. *International Journal of Obesity, 24*, 600-606.
- Schoeller, A. D. (1998). Balancing energy expenditure and body weight. *American Journal of Clinical Nutrition, 68*, 956S-961S.
- Schoeller, A. D., Shay, K., & Kushner, F. R. (1997). How much physical activity is needed to minimize weight gain in previously obese women? *American Journal of Clinical Nutrition, 66*, 551-556.
- Tate, F. D., Jeffery, W. R., Sherwood, E. N., & Wing, R. W. (2007). Long-term weight losses associated with prescription of higher physical activity goals. Are higher levels of physical activity protective against weight loss? *American Journal of Clinical Nutrition, 85*, 954-959.
- Thompson, L. D., Rakow, J., & Perdue, M. S. (2004). Relationship between accumulated

- walking and body composition in middle-aged women. *Medicine & Science in Sports & Exercise*, 36, 911-914.
- Treath, S. M., Hou, N., Young, D., & Maynard L. M. (2005). Accelerometry-measured activity on sedentary time and overweight in rural boys and girls. *Obesity Research*, 13, 1606-1614.
- Tremblay, A. Despres, J-P., Leblanc, C., Lynn, C., Ferris, B., Stephens, T., Bouchard, C. (1990). Effect of intensity of physical activity on body fatness and fat distribution. *American Journal of Clinical Nutrition*, 51, 153-157.
- Tudor-Locke, C., Ainsworth, E. B., Thompson, W. R., Addy, L. C., & Jones, A. D. (2001). The relationship between pedometer-determined ambulatory activity and body composition variables. *International Journal of Obesity*, 25, 1571-1578.
- Warnold, I., Carlgren, G., & Krotkiewski, M. (1978). Energy expenditure and body composition during weight reduction in hyperplastic obese women. *American Journal of Clinical Nutrition*, 31, 750-763.
- Weir, T. L., Ayers, W. G., Jackson, S. A., Rossum, C. A., Poston, C., & Foreyt, P. J. (2001). Determining the amount of physical activity needed for long-term weight control. *International Journal of Obesity*, 25, 613-621.
- Yanai, A., Kon, A., Kumasaka, K., & Kawano, K. (1997). Body mass index variations by age and sex, and prevalence of overweight in Japanese adults. *International Journal of Obesity*, 21, 484-488.